

Genomic DNA from *Anaplasma phagocytophilum*, Strain NCH-1

Catalog No. NR-51150

For research use only. Not for use in humans.

Contributor:

Jason A. Carlyon, Ph.D., Department of Microbiology and Immunology, Virginia Commonwealth University, Richmond, Virginia, USA

Manufacturer:

BEI Resources

Product Description:

Genomic DNA was extracted from a preparation of cell lysate and supernatant from human promyelocytic leukemia cells (HL-60; ATCC® CCL-240™) infected with *Anaplasma phagocytophilum* (*A. phagocytophilum*), strain NCH-1. *A. phagocytophilum*, strain NCH-1 was isolated in 1997 from the blood of a patient suffering from human granulocytic ehrlichiosis in Massachusetts, USA.¹ The complete genome of *A. phagocytophilum*, strain NCH-1 has been sequenced (GenBank: [LANT01000009](#)).

A. phagocytophilum is an intracytoplasmic bacterium that is transmitted by *Ixodes persulcatus* complex ticks and infects bone marrow-derived mammalian cells. It is the causative agent of tick-borne fever of ruminants, equine granulocytic ehrlichiosis, and human granulocytic ehrlichiosis (HGE).^{2,3} *A. phagocytophilum* was formerly known as *Ehrlichia phagocytophila*, but was subsequently reassigned to the *Anaplasmataceae* family.³

NR-51150 has been qualified for PCR applications by amplification of approximately 1470 base pairs of the 16S ribosomal RNA gene.

Material Provided:

Each vial contains approximately 100 µL of bacterial genomic DNA in TE buffer (10 mM Tris-HCl, 1 mM EDTA). The bacterial genomic DNA is in a background of cellular nucleic acid. The vial should be centrifuged prior to opening.

Packaging/Storage:

NR-51150 was packaged aseptically in screw-capped plastic cryovials. The product is provided frozen on dry ice and should be stored at -20°C or colder immediately upon arrival. Freeze-thaw cycles should be minimized.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: Genomic DNA from *Anaplasma phagocytophilum*, strain NCH-1, NR-51150."

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. [Biosafety in Microbiological and Biomedical Laboratories \(BMBL\)](#), 6th ed. Washington, DC: U.S. Government Printing Office, 2020.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Telford, S. R., et al. "Perpetuation of the Agent of Human Granulocytic Ehrlichiosis in a Deer Tick-Rodent Cycle." *Proc. Natl. Acad. Sci. USA* 93 (1996): 6209-6214. PubMed: 8650245.
2. Dumler, J. S., et al. "Human Granulocytic Anaplasmosis and *Anaplasma phagocytophilum*." *Emerg. Infect. Dis.* 11 (2005): 1828-1834. PubMed: 16485466.
3. Dumler, J. S., et al. "Reorganization of Genera in the Families *Rickettsiaceae* and *Anaplasmataceae* in the Order *Rickettsiales*: Unification of Some Species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, Descriptions of Six New

Species Combinations and Designation of *Ehrlichia equi* and 'HGE agent' as Subjective Synonyms of *Ehrlichia phagocytophila*." Int. J. Syst. Evol. Microbiol. 51 (2001): 2145-2165. PubMed: 11760958.

ATCC® is a trademark of the American Type Culture Collection.

